Krusenstern, I. F. 1814. Atlas zur Reise um die Welt unternommen auf Befehl seiner Kaiserlichen Majestät Alexander des Ersten auf den Schiffen Nadeshda und Neva unter dem Commando des Capitans von Krusenstern. St Petersburg.

MacLean, S. F., Jr & Holmes, R. T. 1971. Bill lengths, wintering areas, and taxonomy of

North American Dunlins, Calidris alpina. Auk 88: 893-901.

Meininger, P. L. & van Swelm, N. D. 1989. Steltlopers in de Oosterschelde, biometrisch en ringonderzoek in voorjaar 1984 en 1985. Rijkswaterstaat Dienst Gefijdewateren nota GWAO-89.1009, Stichting Ornithologisch Station Voorn.

Morrison, R. I. G. 1984. Migration systems of some new world shorebirds. Pp. 125-202 in J. Burger & B. L. Olla (eds), Shorebirds Migration and Foraging Behavior (Behavior

Marine Animals, Vol. 6.) Plenum Press.

Nechaev, V. A. 1978. [New data on distribution and biology of birds at Sakhalin Island.]

Ornitologiya 14: 108-111. (In Russian.)

Nechaev, V. A. & Tomkovich, P. S. 1987. [A new subspecies of the Dunlin, Calidris alpina litoralis ssp. n. (Charadriidae, Aves), from the Sakhalin Island.] Zool. Zhurnal 66: 1110-1113. (In Russian.)

Nechaev, V. A. & Tomkovich, P. S. 1988. [A new name for Sakhalin Dunlin (Aves,

Charadriidae).] Zool. Zhurnal 67(10): 1596. (In Russian.)

Norton, D. W. 1971, Two Soviet recoveries of Dunlins banded at Pt. Barrow, Alaska, Auk 88: 927.

Portenko, L. A. 1972. Birds of the Chukchi Peninsula and Wrangel Island. Vol. 1. Acad. Sci. U.S.S.R., Inst. Zool. (English translation, Smithsonian Inst. and Nat. Sci. Found.)

Prater, A. J., Marchant, J. H. & Vuorinen, J. 1977. Guide to the Identification and Ageing of Holarctic Waders. Field Guide no. 17, British Trust for Ornithology.

Sauer, E. G. F. & Urban, E. K. 1964. Bird notes from St Lawrence Island, Alaska. Bonn.

Zool. Beitr. 15: 45-58. Sealy, S. G., Bedard, J., Fay, F. H. & Udvardy, M. D. F. 1971. New records and zoogeographical notes on the birds of St Lawrence Island, Bering Sea. Condor 73: 322–336.

Svetovidov, A. N. 1981. The Pallas fish collection and the Zoographia Rosso-Asiatica: an historical account. Archives of Natural History 10: 45-64.

Thompson, M. C. & DeLong, R. L. 1969. Birds new to North America and the Pribilof Islands, Alaska. Auk 86: 747-749.

Tomkovich, P. S. 1986. [Geographical variability of the Dunlin in the Far East.] Byill. Mosk. Obshch. Ispyt. Prir. 9: 3-15. (In Russian.)

Vaurie, C. 1965. The Birds of the Palearctic Fauna. Non-Passeriformes. H. F. and G. Witherby.

Vieillot, L. J. P. 1816. Nouveau Dictionnaire d'Histoire Naturelle. New ed. Vol. 3. Paris.

Address: M. Ralph Browning, Biological Survey, National Ecology Research Center, U.S. Fish and Wildlife Service, National Museum of Natural History, Washington DC. 20560, U.S.A.

© British Ornithologists Club 1991

Discovery of the nest of Humblot's Flycatcher Humblotia flavirostris

by M. Herremans, M. Louette & 7. Stevens

Received 28 November 1990

Humblot's Flycatcher forms a monotypic genus, probably close to Muscicapa (Benson 1960). The species is confined to a few thousand hectares of mountain forest on the slopes of the Karthala volcano (Louette 1988), which dominates the southern half of Grand Comoro island. Although confiding and rather common, nothing is known as yet about the species' breeding habits, despite the breeding grounds having been

well explored on some occasions (1965, 1983, 1985) during the presumed breeding period (October-November). Benson (1960) speculated that the finding of a nest would be of particular interest in establishing the

affinities of this peculiar bird.

On 12 November 1989, when carrying out point-transect counts in the Karthala forest just north of M'Lima Manda (1050 m a.s.l.) on behalf of an ICBP bird inventory project, M.H. observed a pair of *Humblotia* with food. It turned out that he was standing right under an active nest. The surrounding habitat was a low, rather open type of pioneer woodland on the southern of the two lava flows to the north of M'Lima Mani. The canopy height was only about 12-15 m, and the nest was situated in the mid stratum. The nesting tree had a stem of about 15 cm diameter, was about 8 m high and had a crown with rather dense, stiff, leathery foliage. The nest was in the upper branches, and hidden among the foliage. At least two young were begging. Food items brought to the nest were seen at close range on five occasions. Four of these were single, large centipedes (Chilopoda), and in the remaining case the bird carried two items, one of which was long-legged and presumably a mantis or large spider.

The spot was revisited on 17 December in order to collect the nest. No droppings or feather sheaths remained in it, making it questionable that the young fledged successfully. The nest was an untidy cup or bowl of mosses, lined on the inside with dry grasses, simply resting on (not attached to) some smaller branches of the tree-top (not a fork). The internal diameter of the cup was 5.5 cm and its depth 4 cm. Externally the nest was 7 cm deep, and its base was 15 cm long and 12 cm wide. It was collected and is now stored in the KMMA Museum, Tervuren, Belgium

(nr. 90-01-A-1).

Is there information for the evaluation of the affinities of *Humblotia* in the position of the nest, its simple shape or its constituent materials? These features are rather diverse in African Muscicapa (Erard 1990), but among the species studied in Gabon, in its site and construction the nest of Humblotia comes close to that described for Muscicapa sethsmithi, also a forest species. Based on information on the nest alone, there is apparently no strong reason for sustaining the genus Humblotia apart from Muscicapa.

Acknowledgements

The authors' birding activities on Grand Comoro were sponsored by the ICBP project 22/KM/3. D. Tonnoeyr is thanked for help and for putting on a brave face during the second strenuous walk to M'Lima Manda.

References:

Benson, C. W. 1960. The birds of the Comoro Islands: results of the BOU centenary Expedition 1958. Ibis 103B: 55-58.

Erard, C. 1990. Ecologie et comportement des gobe-mouches (Aves: Muscicapinae, Platysteirinae, Monarchinae) du Nord-Est du Gabon. Vol. 2: Organisation sociale et écologie de la reproduction des Muscicapinae, Mém. Mus. natn. Hist. nat. (A) No. 146, Paris. Louette, M. 1988. Les oiseaux des Comores. Ann. Mus. r. Afr. centr., Serie In-8° No 255,

Tervuren.

Address for correspondence: Dr M. Louette, Koninklijk Museum Midden Afrika, 3080 Tervuren, Belgium.

© British Ornithologists Club 1991